

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
13 September 2001 (13.09.2001)

PCT

(10) International Publication Number
WO 01/66155 A2(51) International Patent Classification⁷: A61K 51/04,
51/10Jackson, TX 77566 (US). GULYAS, Gyongyi [US/US];
305 Timbercreek Drive, Lake Jackson, TX 77566 (US).

(21) International Application Number: PCT/US01/05927

(74) Agent: KARADZIC, Dragan, J.; The Dow Chemical
Company, Intellectual Property, B-1211, 2301 N. Bra-
zosport Boulevard, Freeport, TX 77541 (US).

(22) International Filing Date: 23 February 2001 (23.02.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/185,220 25 February 2000 (25.02.2000) US(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CZ, DE,
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,
ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT,
LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ,
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT,
TZ, UA, UG, US, UZ, YU, ZA, ZW.

(71) Applicant and

(72) Inventor: MA, Dangshe [US/US]; 13011 Twinbrook
Parkway, Rockville, MD 20851 (US).(84) Designated States (*regional*): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): MCDEVITT,
Michael, R. [US/US]; Apartment 6A, 5644 Netherland
Avenue, Bronx, NY 10471 (US). SCHEINBERG, David,
A. [US/US]; 325 Central Park West, New York, NY
10025 (US). SIMON, Jaime [US/US]; Route 1, Box
199-A, Angleton, TX 77515 (US). KIEFER, Garry, E.
[US/US]; 114 Juniper Street, Lake Jackson, TX 77566
(US). FRANK, R., Keith [US/US]; 213 Pansy Path, Lake**Published:**— without international search report and to be republished
upon receipt of that reportFor two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: ACTINIUM-225 COMPLEXES AND CONJUGATES FOR RADIOIMMUNOTHERAPY

(57) **Abstract:** Actinium-225 (²²⁵Ac) complexes with functionalized chelants of the formula (I) wherein: each Q is independently hydrogen or (CHR⁵)_pCO₂R; Q¹ is hydrogen or (CHR⁵)_wCO₂R; each R independently is hydrogen, benzyl or C₁-C₄ alkyl; with the proviso that at least two of the sum of Q and Q¹ must be other than hydrogen; each R⁵ independently is hydrogen; C₁-C₄ alkyl or (C₁-C₂ alkyl)phenyl; X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond; n is 0 or 1; m is an integer from 0 to 10 inclusive; p is 1 or 2; r is 0 or 1; w is 0 or 1; with the proviso that n is only 1 when X and/or Y form an additional carbon to carbon bond, and the sum of r and w is 0 or 1; L is a linker/spacer group covalently bonded to, and replaces one hydrogen atom of one of the carbon atoms to which it is joined, said linker/spacer group being represented by the formula (1) wherein s is an integer of 0 or 1; t is an integer of 0 to 20 inclusive; R¹ is an electrophilic or nucleophilic moiety which allows for covalent attachment to an antibody or fragment thereof, or synthetic linker which can be attached to an antibody or fragment thereof, or precursor thereof; and Cyc represents a cyclic aliphatic moiety, aromatic moiety, aliphatic heterocyclic moiety, or aromatic heterocyclic moiety, each of said moieties optionally substituted with one or more groups which do not interfere with binding to an antibody or antibody fragment; with the proviso that when s, t, m, r, and n are 0, then R¹ is other than carboxyl; their pharmaceutically acceptable salts, their conjugates and the use thereof for radioimmunotherapy is disclosed.



WO 01/66155 A2

ACTINIUM-225 COMPLEXES AND CONJUGATES FOR
RADIOIMMUNOTHERAPY

FIELD OF THE INVENTION

5

This invention relates to actinium-225 (^{225}Ac) complexes with functionalized chelants, their conjugates and their use for radioimmunotherapy.

10

BACKGROUND OF THE INVENTION

The use of radionuclides complexed with suitable chelants, as well as their conjugates (that is, such complexes covalently attached to a biologically active carrier, for example, protein) for diagnosis of cancer and/or therapeutic treatment of cancer in mammals is known. These biochemically engineered molecules provide the tumor specificity and the radioisotope provides potent cytotoxicity. See, for example, U.S. Patent Nos. 4,897,254; 5,342,925; 5,435,990; 5,652,361; 5,696,239; and 5,756,065.

It has been recognized that antibody-targeted alpha particles would allow extraordinary potent, single cell-specific killing with minimal toxicity to normal cells or the patient. The use of alpha particles as an alternative to more traditional classes of radiation is derived from the particle's kinetic characteristics and the radioactive half-life of their source isotope, as well as from the properties of the target-selective carrier moiety for the source isotope. The use of alpha emitting radionuclides is highly desirable for the following reasons: (a) a single atom can kill a cell making them hundreds to thousands of times more potent than even the most potent toxins or drugs; (b) the range of alpha particles is only about 50 microns, so that adjacent tissues are not harmed; (c) the chelated atoms on humanized antibodies are

- unlikely to be immunogenic and can be repeatedly dosed;
(d) the radioactive atoms decay to harmless stable atoms;
(e) killing can occur from inside or outside of the cell;
(f) killing is by apoptosis and by double stranded DNA
5 breaks and repair is not likely.

Specific cytotoxic effects of "alpha particle-emitting radioimmunoconjugates" have been demonstrated in several experimental systems. Specific *in vitro* cell-
10 killing has been demonstrated against a human epidermoid cell line using ^{213}Bi - and ^{225}Ac -containing immunoconjugates, see, for example, Kaspersen et al, *Nuclear Medicine Communications*, Vol. 15, pp. 468-476 (1995). Efficient and specific cell kill by the ^{212}Bi -
15 labeled anti-Tac (CD25) monoclonal antibody has been demonstrated against an adult T-cell leukemia cell line *in vitro*, see, for example, R. W. Kozak et al, *Proc. Natl. Acad. Sci. USA*, Vol. 83, pp. 474-478 (1986). In other experiments, mice inoculated intraperitoneally with the
20 murine tumor line EL-4 were cured of their ascites after intraperitoneal injection of 150 μCi of a ^{212}Bi -labeled antibody conjugate, see, for example, R. M. Macklis et al, *Science*, Vol. 240, pp. 1024-1026 (1988).

25 Potential for use of ^{225}Ac in radiotherapy of cancer has also been recognized due to its favorable properties. This isotope decays with a radioactive half-life of 10 days into a cascade of short-lived alpha and beta-emitting isotopes. See, for example, M. W. Geerlings et al,
30 *Nuclear Medicine Communications*, Vol. 14, pp. 121-125 (1993) and Kaspersen et al, *Nuclear Medicine Communications*, Vol. 15, pp. 468-476 (1995). However, the use of ^{225}Ac in radioimmunotherapy has been hampered due to its toxicity and lack of a suitable carrier which will
35 deliver it to the targeted cells.

In an effort to reduce the toxicity of ^{225}Ac , numerous chelating agents such as, for example, 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetraacetic acid (DOTA), diethylenetriaminepentaacetic acid (DTPA), ethylene-
5 diaminetetracetic acid (EDTA), 1,4,7,10,13-pentaazacyclopentadecane-1,4,7,10,13-pentaacetic acid (PEPA), and 1,4,7,10,13,16-hexaazacyclohexadecane-1,4,7,10,13,16-hexaacetic acid (HEHA) have been complexed with ^{225}Ac and evaluated *in vivo* for toxicity and stability. However,
10 the toxicity of these complexes has proved to be still substantial.

G. J. Beyer et al, *Isotoperpraxis*, Vol. 26, pp. 111-114 (1990), has evaluated the *in vivo* uptake of ^{225}Ac -
15 citrate and compared it to ^{169}Yb -citrate. This study has found that ^{225}Ac -citrate had more efficient blood clearance, greater liver uptake, and lower bone uptake than ^{169}Yb -citrate.

G. J. Beyer et al, *Nucl. Med. & Biol.*, Vol. 24, pp. 367-372 (1997), has evaluated EDTMP (ethylenediaminetetramethylenephosphonic acid) as a chelant for ^{225}Ac . The
20 study has found that EDTMP, depending on its concentration, reduces the liver uptake. However, the liver uptake of ^{225}Ac -EDTMP is still substantial and
25 excretion of ^{225}Ac -EDTMP is poor. The study has also suggested that greater efficacy in endoradionuclide therapy of bone metastasis can be expected with the use of ^{225}Ac -EDTMP due to the alpha-radiation.

30 K. A. Deal et al, *J. Med. Chem.*, Vol 42, pp. 298-2992 (1999), has evaluated biodistribution of a number of ^{225}Ac chelates. It has been observed that the structure of the chelant has dramatic effect on biodistribution of ^{225}Ac .
35 HEHA (1,4,7,10,13,16-hexaazacyclohexadecane-1,4,7,10,13,16-hexaacetic acid) was the largest

macrocyclic chelant. ^{225}Ac readily formed a complex with HEHA. Exceptional in vivo stability and reduced toxicity has been observed for ^{225}Ac -HEHA. This has been attributed to the large size and macrocyclic effect of HEHA.

5

Although various chelating agents were suggested and evaluated as carriers for ^{225}Ac , up to now ^{225}Ac has not been successfully chelated to an antibody and no successful therapeutic use of ^{225}Ac in animals or humans has been reported presumably due to its inherent toxicity and/or stability problems of its complexes.

It would be desirable to provide complexes comprising ^{225}Ac and functionalized chelants which are kinetically and thermodynamically inert for use in therapeutic applications.

It would also be desirable to provide conjugates of such ^{225}Ac complexes with a biological molecule. The biological molecule in these conjugates would provide the tumor specificity and the ^{225}Ac isotope would provide potent cytotoxicity.

Another desirable property of these conjugates includes physiological compatibility which would permit the ^{225}Ac complex, if separated from its targeting, conjugated biological molecule *in vivo*, to be soluble in physiological fluids and thus be rapidly eliminated from the body.

30


SUMMARY OF THE INVENTION

The present invention is directed to ^{225}Ac complexes and their conjugates with a biological molecule. The ^{225}Ac complexes and conjugates of the present invention are

35

useful for the treatment of cancer in mammals, especially humans.


More specifically, the present invention is directed to ²²⁵Ac complexes comprising a functionalized chelant compound of the formula (I):



wherein:

- each Q is independently hydrogen or (CHR⁵)_pCO₂R;
- Q¹ is hydrogen or (CHR⁵)_wCO₂R;
- each R independently is hydrogen, benzyl or C₁-C₄ alkyl; with the proviso that at least two of the sum of Q and Q¹ must be other than hydrogen;
- each R⁵ independently is hydrogen; C₁-C₄ alkyl or (C₁-C₂ alkyl)phenyl;
- X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;
- n is 0 or 1;
- m is an integer from 0 to 10 inclusive;
- p is 1 or 2;
- r is 0 or 1;
- w is 0 or 1;
- with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of r and w is 0 or 1;
- L is a linker/spacer group covalently bonded to, and replaces one hydrogen atom of one of the carbon atoms

to which it is joined, said linker/spacer group being represented by the formula



5

wherein

s is an integer of 0 or 1;

t is an integer of 0 to 20 inclusive;

10 R^1 is an electrophilic or nucleophilic moiety which allows for covalent attachment to an antibody or fragment of thereof, or synthetic linker which can be attached to an antibody or fragment thereof, or precursor thereof; and

15 Cyc represents a cyclic aliphatic moiety, aromatic moiety, aliphatic heterocyclic moiety, or aromatic heterocyclic moiety, each of said moieties optionally substituted with one or more groups which do not interfere with binding to a biologically active carrier;

20

with the proviso that when s, t, m, r, and n are 0, then R^1 is other than carboxyl; or pharmaceutically acceptable salt thereof; complexed with ^{225}Ac .

25

The present invention is also directed to a conjugate comprising the aforementioned ^{225}Ac complex covalently attached to a biological molecule.

30

The present invention also includes formulations having the conjugates of this invention and a pharmaceutically acceptable carrier, especially

formulations where the pharmaceutically acceptable carrier is a liquid.

5 The present invention is also directed to a method of therapeutic treatment of a mammal having cancer which comprises administering to said mammal a therapeutically effective amount of the formulation of this invention.

10 Surprisingly, the ^{225}Ac complexes and conjugates of this invention are relatively stable (that is, do not easily dissociate) and some display rapid clearance from the whole body and some non-target organs, such as liver and kidney. Additionally, the alpha-particles emitting ^{225}Ac complexes and conjugates of this invention are
15 expected to have several advantages over beta particle-emitting cytotoxic agents including higher energy and more potent emissions, less hazardous waste, expected lower effective dose, the potential for outpatient treatment, better retention at the target sites, and higher target to
20 non-target radiation ratios.

DETAILED DESCRIPTION OF THE INVENTION

25 As used herein, the term " ^{225}Ac complex" refers to a functionalized chelant compound of formula I complexed with ^{225}Ac radionuclide.

30 As used herein, the term " ^{225}Ac conjugate" refers to ^{225}Ac complex of the present invention that is covalently attached to a biological molecule.

As used herein, the term "mammal" means animals that
nurish their young with milk secreted by mammary glands,
preferably humans.

35

As used herein, the term "biological molecule" refers to any protein, antibody, antibody fragment, hormone, peptide, growth factor, antigen, hapten or any other carrier which functions in this invention to recognize a specific biological target site. Antibody and antibody fragment refers to any polyclonal, monoclonal, chimeric, human, mammalian, single chains, dimeric and tetrameric antibody or antibody fragment. Such biological molecule, when attached to a functionalized complex, serves to carry the attached ^{225}Ac ion to specific targeted tissues. The term "antibody" refers to any polyclonal, monoclonal, chimeric antibody or heteroantibody. Preferably the antibodies used in the ^{225}Ac conjugates of the present invention are monoclonal antibodies having high specificity for the desired cancer cells. Antibodies used in the present invention may be directed against, for example, cancer, tumors, leukemias, autoimmune disorders involving cells of the immune system, normal cells that need to be ablated such as bone marrow and prostate tissue, virus infected cells including HIV, mycoplasma, differentiation and other cell membrane antigens, pathogen surface antigens and any biologically active molecules. Some examples of antibodies are HuM195 (anti-CD33), CC-11, CC-46, CC-49, CC-49 F(ab')₂, CC-83, CC-83 F(ab')₂, and B72.3. Particularly preferred antibody for use in the practice of the present invention is HuM195. Antibody fragment includes Fab fragments and F(ab')₂ fragments, and any portion of an antibody having specificity toward a desired epitope or epitopes. The antibodies which may be used in the ^{225}Ac conjugates of the present invention can be prepared by techniques well known in the art. Highly specific monoclonal antibodies can be produced by hybridization techniques well known in the art, see, for example, Kohler and Milstein, *Nature*, 256, 495-497 (1975); and *Eur. J. Immunol.*, 511-519 (1976).

As used herein, "pharmaceutically acceptable salt" means any salt of a compound of formula (I) which is sufficiently non-toxic to be useful in therapy of mammals. Representative of those salts, which are formed by standard reactions, from both organic and inorganic sources include, for example, sulfuric, hydrochloric, phosphoric, acetic, succinic, citric, lactic, maleic, fumaric, palmitic, cholic, palmoic, mucic, glutamic, d-camphoric, glutaric, glycolic, phthalic, tartaric, formic, lauric, steric, salicylic, methanesulfonic, benzenesulfonic, sorbic, picric, benzoic, cinnamic and other suitable acids. Also included are salts formed by standard reactions from both organic and inorganic sources such as ammonium, alkali metal ions, alkaline earth metal ions, and other similar ions. Preferred are the salts of the compounds of formula I where the salt is potassium, sodium, ammonium, or mixtures thereof.

As used herein, the term "therapeutically effective amount" means an amount of the ^{225}Ac conjugate that produces a therapeutic effect on the disease treated. The therapeutically effective amount will vary depending on the mammal, the ^{225}Ac conjugate and the method of its administration (for example, oral or parenteral). A person of ordinary skill in the art can determine the therapeutically effective amount of the ^{225}Ac conjugate.

In the practice of the present invention the ^{225}Ac conjugate may be administered *per se* or as a component of a pharmaceutically acceptable formulation.

Thus, the present invention may be practiced with the ^{225}Ac conjugate being provided in pharmaceutical formulation, both for veterinary and for human medical use. Such pharmaceutical formulations comprise the active agent (the ^{225}Ac conjugate) together with a physiologically

acceptable carrier, excipient or vehicle therefore. The carrier(s) must be physiologically acceptable in the sense of being compatible with the other ingredient(s) in the formulation and not unsuitably deleterious to the recipient thereof. The ^{225}Ac conjugate is provided in a therapeutically effective amount, as described above, and in a quantity appropriate to achieve the desired dose.

The formulations include those suitable for parenteral (including subcutaneous, intramuscular, intraperitoneal, and intravenous), oral, rectal, topical, nasal, or ophthalmic administration. Formulations may be prepared by any methods well known in the art of pharmacy. Such methods include the step of bringing the ^{225}Ac conjugate into association with a carrier, excipient or vehicle therefore. In general, the formulation may be prepared by uniformly and intimately bringing the ^{225}Ac conjugate into association with a liquid carrier, a finely divided solid carrier, or both, and then, if necessary, shaping the product into desired formulation. In addition, the formulations of this invention may further include one or more accessory ingredient(s) selected from diluents, buffers, binders, disintegrants, surface active agents, thickeners, lubricants, preservatives, and the like. In addition, a treatment regime might include pretreatment with non-radioactive carrier.

Injectable formulations of the present invention may be either in suspensions or solution form. In the preparation of suitable formulations it will be recognized that, in general, the water solubility of the salt is greater than the acid form. In solution form the complex (or when desired the separate components) is dissolved in a physiologically acceptable carrier. Such carriers comprise a suitable solvent, preservatives such as free radical quenching agents, for example, ascorbic acid,

benzyl alcohol or any other suitable molecule, if needed, and buffers. Useful solvents include, for example, water, aqueous alcohols, glycols, and phosphonate or carbonate esters. Such aqueous solutions contain no more than 50 percent of the organic solvent by volume.


Injectable suspensions are compositions of the present invention that require a liquid suspending medium, with or without adjuvants, as a carrier. The suspending medium can be, for example, aqueous polyvinylpyrrolidone, inert oils such as vegetable oils or highly refined mineral oils, polyols, or aqueous carboxymethylcellulose. Suitable physiologically acceptable adjuvants, if necessary to keep the complex in suspension, may be chosen from among thickeners such as carboxymethylcellulose, polyvinylpyrrolidone, gelatin, and the alginates. Many surfactants are also useful as suspending agents, for example, lecithin, alkylphenol, polyethyleneoxide adducts, naphthalenesulfonates, alkylbenzenesulfonates, and polyoxyethylene sorbitane esters.

In the context of the present invention the terms "functionalized chelant" and "bifunctional chelant" are used interchangeably and refer to compounds which have the dual functionality of sequestering metal ions plus the ability to covalently bind a biological molecule having specificity for tumor cell epitopes or antigens. Such compounds are of great utility for therapeutic and diagnostic applications when they are, for example, complexed with radioactive metal ions and covalently attached to a specific antibody. These types of complexes have been used to carry radioactive metals to tumor cells which are targeted by the specificity of the attached antibody [see, for example, Mears et al., *Anal. Biochem.*

142, 68-74 (1984); Krejcarek et al., Biochem. And Biophys. Res. Comm. 77, 581-585 (1977)].

The functionalized chelant compounds of formula (I)
 5 useful in the practice of the present invention are known in the art. See, for example, U.S. Patent Nos. 5,435,990 and 5,652,361.

Compounds of formula I where: R is hydrogen or
 10 methyl; n is 0; m is 0 through 5; r is 0; and L is a moiety of formula A:



15 wherein:

R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;


20 R^3 is selected from the group consisting of C_1 - C_4 alkoxy, $-OCH_2CO_2H$, hydroxy and hydrogen; and

R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and
 25 maleimido;

with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof; are preferred functionalized chelants.

30


Preferred functionalized chelant compounds of formula I include also those compounds where Q^1 is hydrogen and L is represented by formula A as shown by formula II:



wherein:

- 5 each Q independently is hydrogen or CHR^5COOR ; with the proviso that at least two of Q must be other than hydrogen
- each R independently is hydrogen benzyl or $\text{C}_1\text{-C}_4$ alkyl;
- 10 m is integer from 0 to 5 inclusive;
- R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 15 R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$ alkoxy, $-\text{OCH}_2\text{COOH}$, hydroxy and hydrogen;
- R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 20 each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl; with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen;
- or
- 25 a pharmaceutically acceptable salt thereof.


Additional preferred functionalized chelant compounds of formula I include those compounds where at least one Q is hydrogen and are represented by formula III:



wherein:

- 5 each Q independently is hydrogen or CHR^5COOR ;
 Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$; with the proviso that at
 least two the sum of Q and Q^1 must be other than
 hydrogen and one Q is hydrogen;
 each R independently is hydrogen benzyl or $\text{C}_1\text{-C}_4$
 10 alkyl;
 m is integer from 0 to 5 inclusive;
 w is 0 or 1;
 R^2 is selected from the group consisting of hydrogen,
 nitro, amino, isothiocyanato, semicarbazido,
 15 thiosemicarbazido, carboxyl, bromoacetamido and
 maleimido;
 R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$
 alkoxy, $-\text{OCH}_2\text{COOH}$, hydroxy and hydrogen;
 R^4 is selected from the group consisting of hydrogen,
 20 nitro, amino, isothiocyanato, semicarbazido,
 thiosemicarbazido, carboxyl, bromoacetamido and
 maleimido;
 each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl;
 with the proviso that R^2 and R^4 cannot both be
 25 hydrogen but one of R^2 and R^4 must be hydrogen; or
 a pharmaceutically acceptable salt thereof.

Other preferred functionalized chelant compounds of
 formula I include compounds where Q^1 is CO_2R ($w=0$) and are
 30 represented by formula IV:



wherein:

- 5 each Q independently is hydrogen or CHR⁵COOR; with the proviso that at least one Q must be other than hydrogen;
- each R independently is hydrogen benzyl or C₁-C₄ alkyl;
- 10 m is integer from 0 to 5 inclusive;
- R² is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 15 R³ is selected from the group consisting of C₁-C₄ alkoxy, -OCH₂COOH, hydroxy and hydrogen;
- R⁴ is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 20 each R⁵ independently is hydrogen or C₁-C₄ alkyl; with the proviso that R² and R⁴ cannot both be hydrogen but one of R² and R⁴ must be hydrogen; or a pharmaceutically acceptable salt thereof.

25

 The functionalized chelants of formula I useful in the practice of the present invention can be prepared by known methods. General synthetic approach to a twelve-membered macrocyclic, bifunctional chelant of the present invention as represented by formula I involves

30 monofunctionalization of a free-base macrocycle (for

example, 1,4,7,10-tetraazacyclododecane) at only one of the nitrogen atoms with an appropriate electrophile (for example, any appropriately substituted alpha-halocarboxylic acid ester). This electrophile must
5 possess a suitable linker moiety which would allow covalent attachment of bifunctional ligand to a biological molecule. Various synthetic routes to functionalized chelants of formula I have been described U.S. Patent Nos. 5,435,990 and 5,652,361, both incorporated herein by
10 reference.

The method of obtaining ^{225}Ac radionuclide is not critical to the present invention. For example, ^{225}Ac can be prepared in a cyclotron. ^{225}Ac can be obtained in pure
15 form from Department of Energy (DOE), U.S.A., and Institute for Transuranium Elements (ITU), Karlsruhe, Germany.

The ^{225}Ac conjugates of the present invention can be
20 prepared by first forming the complex and then binding the biological molecule. Thus, the process involves preparing or obtaining the ligand, forming the complex with ^{225}Ac and then adding the biological molecule. Alternatively, the process may involve first conjugation of the ligand to the
25 biological molecule and then the formation of the complex with ^{225}Ac . Any suitable process that results in the formation of the ^{225}Ac conjugates of this invention is within the scope of the present invention.

30 In the following examples, the following terms and conditions were used unless otherwise specified.

Glossary of Terms

35 Ab = antibody;
BFC = bifunctional chelant;

DOTA = 1,4,7,10 tetraazacyclododecane-1,4,7,10-tetraacetic acid;
MeO-DOTA-NCS = 1-[(2-methoxy-5-isothiocyanato-phenyl)-carboxymethyl]-4,7,10-triscarboxy-
5 methyl-1,4,7,10-tetraazacyclododecane;
TMAA = tetramethyl ammonium acetate buffer;
Sephadex C-25 resin is a cation exchange resin, sold by Pharmacia Inc.;
EDTA = ethylenediaminetetraacetic acid;
10 DTPA = diethylenetriaminepentaacetic acid;
TETA = 1,4,8,11-tetraazacyclotetradecane-1,4,8,11-tetraacetic acid;
DOTPA = 1,4,7,10-tetraazacyclododecane-1,4,7,10-tetrapropionic acid;
15 TETPA = 1,4,8,11-tetraazacyclotetradecane-1,4,8,11-tetrapropionic acid;
DOTMP = 1,4,6,10-tetraazacyclodecane-1,4,7,10-tetramethylenephosphonic acid.

20 General Experimental

Method of preparation of ^{225}Ac conjugates: The preparation of ^{225}Ac conjugates involved two steps. First, the ^{225}Ac complex was prepared by mixing a solution of the
25 functionalized chelant compound of formula I with the solution of ^{225}Ac at pH of about 5-6 in a suitable buffer. The complex formation was tested using cation exchange chromatography. Then, the conjugation of the ^{225}Ac complex to a biological molecule (suitably an antibody) was
30 carried out at the pH of about 8.5 in the presence of a suitable buffer. The antibody and the antibody ^{225}Ac complex conjugates were then separated from the unconjugated low molecular weight materials using gel filtration chromatography. The fraction of radioactivity
35 associated with the antibody was then determined.

The γ emission counting was performed using a 3-inch x 3-inch NaI well crystal utilizing the γ emission of ^{225}Ac decay product ^{221}Fr (half-life of 4.8 min.) at 218 KeV. Counting was carried out half an hour after sample preparation.

Method for determining yield and stability of ^{225}Ac complexes and conjugates thereof: Instant Thin Layer Chromatography (ITLC) was utilized with either a 10 mM EDTA or 10 mM NaOH/9% NaCl solvent systems using ITLC SG strips (sold by Gelman Sciences company) to assess the complexation and conjugation efficiency of the DOTA-based bifunctional ^{225}Ac conjugate with HuM195 antibody.

The following examples are provided to further illustrate the present invention, and should not be construed as limiting thereof.

Example 1: Preparation of ^{225}Ac -MeO-DOTA-NCS Complex

An aqueous solution of MeO-DOTA-NCS (35 μl ; 0.31 mg/ml) was mixed with the ^{225}Ac chloride solution (35 μl ; 1.65 $\mu\text{Ci}/\mu\text{l}$,) in 0.1M HCl. The pH was adjusted to about 5 using the TMAA buffer (130 μl , 0.2 M, pH about 6). Reaction mixture was incubated at about 50°C for one hour. Complex formation was checked by cation exchange chromatography employing the Sephadex C-25 resin and it was determined that 99 percent of ^{225}Ac was complexed.

Example 2: Preparation of ^{225}Ac -HuM195 Conjugate

HuM195 antibody solution (20 μl , 5 mg/ml) was added to the ^{225}Ac complex solution (200 μl) prepared as described in Example 1. The pH was adjusted to about 8.5 using a NaHCO_3 buffer (85 μl , 0.1 M, pH=8.7). The molar ratios of the reactants used were as follows: MeO-DOTA-

NCS / ^{225}Ac = 6549; MeO-DOTA-NCS / HuM195 = 24; and HuM195 / ^{225}Ac = 275. After 30 minutes incubation at 20°C the protein and the small molecular weight components of the solution were separated by gel filtration chromatography using the Econo-Pack 10 DG gel filtration column. The extent of coupling was determined by γ emission counting. It was determined that 7.3 percent of the ^{225}Ac complex was coupled to HuM195 antibody.

10 Example 3: Conjugation of Antibodies HuM195 (anti-CD33) and B4 (anti-CD19) to MeO-DOTA-NCS

20 mg of HuM195 (or B4) monoclonal antibody solution was mixed with 0.05 M HEPES containing EDTA at about pH 8 and dialysed against 0.05 M HEPES buffer for 24 hours in an Amicon stirred cell dialysis unit to remove any metal ions associated with antibodies. A MeO-DOTA-NCS solution containing 3.38 mg of MeO-DOTA-NCS was added and allowed to conjugate with the antibody at room temperature for 24 hours. Then the reaction mixture was dialysed against a NaAc / NaCl buffer solution at about pH 7.0 for 24 hours to remove any unreacted MeO-DOTA-NCS. The immunoconjugate was recovered from the stirred cell and characterized using a size exclusion high pressure liquid chromatography (HPLC). This was compared to the native HuM195 or B4. Antibody concentration was determined by UV-absorption at 280 nM.

The immunoconjugates could be labeled readily with ^{111}In showing success of the conjugation reaction. For example, approximately 400 μCi of ^{111}In in 200 μl of 0.2 M HCl was mixed with 29 μl of ammonium acetate (3M) and 9 μl of l-ascorbic acid (150 mg/ml) to adjust the pH to 5.0 and then 0.5 mg of HuM195-MeO-DOTA-NCS immunoconjugate was added. The reaction was allowed to progress at 37°C for 60 minutes. An 87% reaction yield was obtained.

Example 4: Labeling of HuM195-MeO-DOTA
Immunoconjugate with ^{225}Ac

200 μCi of ^{225}Ac in 0.2 M HCl was mixed with 700 μl
5 metal free water and then buffered with 93 μl of NH_4Ac at
about pH 6.5. Each of four 200 μl aliquots was mixed with
0 mg, 0.1 mg, 0.5 mg, 1 mg of HuM195-MeO-DOTA (prepared as
in Example 3), respectively. The reaction tubes were
placed into a 37°C water bath. The ^{225}Ac incorporation was
10 monitored by TLC developed in 10 mM EDTA solvent at 3
hours. The ^{225}Ac incorporation (percentage of activity
remaining at origin) data are given in Table 1 below for
the various antibody concentrations and Ab:Ac ratios.

15

Table 1: ^{225}Ac Incorporation at 3 hr for three different
concentrations of the antibody

Antibody (μM)	1.6	8.1	16.1
Ab to Ac Ratio	174	870	1740
% Incorporation	7.1	23.1	51.8

20

Example 5: Stability of ^{225}Ac -HuM195-MeO-DOTA

The three ^{225}Ac -HuM195-MeO-DOTA solutions from Example
4 were combined and challenged with 20 μl of 10 mM DTPA to
remove unbound metals. 100 μl of L-ascorbic acid (150
25 mg/ml) was added as a radioprotection agent. The solution
was purified through a 10-DG desalting column (Bio-Rad
company) to separate ^{225}Ac -HuM195-MeO-DOTA from unreacted
 ^{225}Ac (in DTPA form). The purified ^{225}Ac -HuM195-MeO-DOTA
was subjected to a stability study in human serum.

30

The purified ^{225}Ac -HuM195-MeO-DOTA was assessed for
stability in different media such as 1% and 25% albumin
(human) and 1% and 25% human serum at 37°C . Overall
stability half-lives of ^{225}Ac -HuM195-MeO-DOTA in either

albumin (human) at 4 °C or human serum at 37 °C exceeded 150 days.

Example 6: ^{225}Ac -HuM195-MeO-DOTA *In-Vitro* Cell Kill

5

A cell-based immunoreactivity study (binding of labeled antibody to antigen excess) has shown that ^{225}Ac labeled HuM195 antibody (^{225}Ac -HuM195-MeO-DOTA) is still immunoreactive (~70%). The potency and specificity of ^{225}Ac -HuM195-MeO-DOTA was then evaluated *in-vitro* as a function of specific activity and activity concentration on antigen positive and negative cell lines. The LD₅₀ in a 5-day assay was ~0.3 nCi/ml at a specific activity of 0.035 Ci/g which is 3 log more potent than the similar ^{213}Bi alpha emitting agent with much high specific activity (see Nikula et al, *J Nucl Med* 1999; 40, 166-176). These data (derived from 3H-thymidine incorporation) are plotted below in Figure 3. The LD₅₀ in a 2-day assay is about 1.4 nCi/ml for positive cell line and about 28 nCi/ml for negative cell line which demonstrates the specificity. Internalization of isotope into target cells was also demonstrated and more than 50% was internalized in the target cells in 5 hours which is crucial to control the fate of daughter isotopes. This preliminary study suggests that ^{225}Ac -HuM195-MeO-DOTA conjugates are useful clinically as a way to target alpha particles to kill cells.

10

15

20

25




Figure 3. ^{225}Ac *In-Vitro* Cell Kill

5

Example 7: *In Vivo* Biodistribution

The *in vivo* biodistribution of free ^{225}Ac acetate, ^{225}Ac -DOTA and ^{225}Ac -HuM195-MeO-DOTA was studied in nu/nu mice by intraperitoneal injection of approximately 2 μCi of ^{225}Ac of each compound in 400 μl . It was demonstrated that a different pattern of distribution existed (see Table 2 below) for the three agents. The ^{225}Ac -DOTA was excreted very quickly and most activity was cleared in less than 40 minutes. ^{225}Ac in acetate was held up in the liver and bone but cleared from blood. The ^{225}Ac -HuM195-MeO-DOTA has longer blood circulation time and less bone uptake over the period of 5 days. These data indicated ^{225}Ac -HuM195-MeO-DOTA is stable *in vivo*.

20


Table 2. Summary of the ^{225}Ac Biodistribution (% dose/g) in nu/nu Mice

	^{225}Ac Acetate			^{225}Ac -DOTA			^{225}Ac -HuM195-DOTA HuM195-MeO-DOTA		
	1 d	2d	5d	40 min	2 h	18 h	1 d	2d	5d
Average									
Blood	0.1	0.2	0.0	3.0	1.8	1.1	10.1	10.2	4.5
Kidneys	3.8	3.2	2.8	4.9	3.5	3.1	4.6	5.2	4.6
Liver	46.6	43.1	68.5	4.5	4.6	5.8	8.8	10.0	20.7
Bone	13.3	12.0	17.1	3.9	4.2	4.9	3.8	4.1	5.3

WHAT IS CLAIMED IS:


1. An ^{225}Ac complex comprising a functionalized chelant compound of the formula I

5



wherein:


- 10 each Q is independently hydrogen or $(\text{CHR}^5)_p\text{CO}_2\text{R}$;
 Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$;
 each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl; with the proviso that at least two of the sum of Q and Q^1 must be other than hydrogen;
 15 each R^5 independently is hydrogen; $\text{C}_1\text{-C}_4$ alkyl or $(\text{C}_1\text{-C}_2 \text{ alkyl})\text{phenyl}$;
 X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;
 20 n is 0 or 1;
 m is an integer from 0 to 10 inclusive;
 p is 1 or 2;
 r is 0 or 1;
 w is 0 or 1;
 25 with the proviso that n is only 1 when X and/or Y form an additional carbon to carbon bond, and the sum of r and w is 0 or 1;
 L is a linker/spacer group covalently bonded to, and replaces one hydrogen atom of one of the carbon atoms to which it is joined, said linker/spacer group being
 30 represented by the formula



wherein

- 5 s is an integer of 0 or 1;
 t is an integer of 0 to 20 inclusive;
 R¹ is an electrophilic or nucleophilic moiety which
 allows for covalent attachment to an antibody or
 fragment of thereof, or synthetic linker which can be
 10 attached to an antibody or fragment thereof, or
 precursor thereof; and
 Cyc represents a cyclic aliphatic moiety, aromatic
 moiety, aliphatic heterocyclic moiety, or aromatic
 heterocyclic moiety, each of said moieties optionally
 15 substituted with one or more groups which do not
 interfere with binding to an antibody or antibody
 fragment;
 with the proviso that when s, t, m, r, and n are 0,
 then R¹ is other than carboxyl; or
 20 a pharmaceutically acceptable salt thereof; complexed
 with ²²⁵Ac.

2. The ²²⁵Ac complex of Claim 1 wherein the
 functionalized chelant is a compound of formula II




wherein:

- 5 each Q independently is hydrogen or CHR⁵COOR; with the proviso that at least two of Q must be other than hydrogen
- each R independently is hydrogen benzyl or C₁-C₄ alkyl;
- 10 m is integer from 0 to 5 inclusive;
- R² is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 15 R³ is selected from the group consisting of C₁-C₄ alkoxy, -OCH₂COOH, hydroxy and hydrogen;
- R⁴ is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 20 each R⁵ independently is hydrogen or C₁-C₄ alkyl; with the proviso that R² and R⁴ cannot both be hydrogen but one of R² and R⁴ must be hydrogen; or a pharmaceutically acceptable salt thereof.

25


3. The ²²⁵Ac complex of Claim 1 wherein the functionalized chelant is a compound of formula III



wherein:

- 5 each Q independently is hydrogen or CHR^5COOR ;
 Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$; with the proviso that at
 least two the sum of Q and Q^1 must be other than
 hydrogen and one Q is hydrogen;
 each R independently is hydrogen benzyl or $\text{C}_1\text{-C}_4$
 10 alkyl;
 m is integer from 0 to 5 inclusive;
 w is 0 or 1;
 R^2 is selected from the group consisting of hydrogen,
 nitro, amino, isothiocyanato, semicarbazido,
 15 thiosemicarbazido, carboxyl, bromoacetamido and
 maleimido;
 R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$
 alkoxy, $-\text{OCH}_2\text{COOH}$, hydroxy and hydrogen;
 R^4 is selected from the group consisting of hydrogen,
 20 nitro, amino, isothiocyanato, semicarbazido,
 thiosemicarbazido, carboxyl, bromoacetamido and
 maleimido;
 each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl;
 with the proviso that R^2 and R^4 cannot both be
 25 hydrogen but one of R^2 and R^4 must be hydrogen; or
 a pharmaceutically acceptable salt thereof.

4. The ^{225}Ac complex of Claim 1 wherein the
 functionalized chelant is a compound of formula IV



wherein:


- 5 each Q independently is hydrogen or CHR^5COOR ; with the proviso that at least one Q must be other than hydrogen;
- each R independently is hydrogen benzyl or $\text{C}_1\text{-C}_4$ alkyl;
- 10 m is integer from 0 to 5 inclusive;
- R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 15 R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$ alkoxy, $-\text{OCH}_2\text{COOH}$, hydroxy and hydrogen;
- R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 20 each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl; with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof.

25

5. The ^{225}Ac complex of Claim 1 wherein the functionalized chelant compound is 1-[(2-methoxy-5-isothiocyanatophenyl)-carboxymethyl]-4,7,10-triscarboxymethyl-1,4,7,10-tetraazacyclododecane (MeO-DOTA-NCS).

30

6. An ^{225}Ac conjugate comprising a functionalized chelant compound of the formula I




5

wherein:

- each Q is independently hydrogen or $(\text{CHR}^5)_p\text{CO}_2\text{R}$;
 Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$;
 10 each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl; with the proviso that at least two of the sum of Q and Q^1 must be other than hydrogen;
 each R^5 independently is hydrogen; $\text{C}_1\text{-C}_4$ alkyl or $(\text{C}_1\text{-C}_2 \text{ alkyl})\text{phenyl}$;
 15 X and Y are each independently hydrogen or may be taken with an adjacent X and Y to form an additional carbon-carbon bond;
 n is 0 or 1;
 m is an integer from 0 to 10 inclusive;
 20 p is 1 or 2;
 r is 0 or 1;
 w is 0 or 1;
 with the proviso that n is only 1 when X and/or Y form an additional carbon-carbon bond, and the sum of
 25 r and w is 0 or 1;
 L is a linker/spacer group covalently bonded to, and replaces one hydrogen atom of one of the carbon atoms to which it is joined, said linker/spacer group being represented by the formula


30



wherein

- 5 s is an integer of 0 or 1;
 t is an integer of 0 to 20 inclusive;
 R¹ is an electrophilic or nucleophilic moiety which
 allows for covalent attachment to an antibody or
 fragment of thereof, or synthetic linker which can be
 10 attached to an antibody or fragment thereof, or
 precursor thereof; and
 Cyc represents a cyclic aliphatic moiety, aromatic
 moiety, aliphatic heterocyclic moiety, or aromatic
 heterocyclic moiety, each of said moieties optionally
 15 substituted with one or more groups which do not
 interfere with binding to an antibody or antibody
 fragment;
 with the proviso that when s, t, m, r, and n are 0,
 then R¹ is other than carboxyl; or
 20 pharmaceutically acceptable salt thereof;
 complexed with ²²⁵Ac; and covalently attached to a
 biological molecule.

7. The ²²⁵Ac conjugate of Claim 6 wherein the
 25 functionalized chelant is a compound of formula II




wherein:

- 5 each Q independently is hydrogen or CHR^5COOR ; with the proviso that at least two of Q must be other than hydrogen
- each R independently is hydrogen benzyl or $\text{C}_1\text{-C}_4$ alkyl;
- 10 m is integer from 0 to 5 inclusive;
- R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 15 R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$ alkoxy, $-\text{OCH}_2\text{COOH}$, hydroxy and hydrogen;
- R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 20 each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl; with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof.

25


8. The ^{225}Ac conjugate of Claim 6 wherein the functionalized chelant is a compound of formula III



wherein:

- 5 each Q independently is hydrogen or CHR^5COOR ;
 Q^1 is hydrogen or $(\text{CHR}^5)_w\text{CO}_2\text{R}$; with the proviso that at
 least two the sum of Q and Q^1 must be other than
 hydrogen and one Q is hydrogen;
 each R independently is hydrogen benzyl or $\text{C}_1\text{-C}_4$
 10 alkyl;
 m is integer from 0 to 5 inclusive;
 w is 0 or 1;
 R^2 is selected from the group consisting of hydrogen,
 nitro, amino, isothiocyanato, semicarbazido,
 15 thiosemicarbazido, carboxyl, bromoacetamido and
 maleimido;
 R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$
 alkoxy, $-\text{OCH}_2\text{COOH}$, hydroxy and hydrogen;
 R^4 is selected from the group consisting of hydrogen,
 20 nitro, amino, isothiocyanato, semicarbazido,
 thiosemicarbazido, carboxyl, bromoacetamido and
 maleimido;
 each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl;
 with the proviso that R^2 and R^4 cannot both be
 25 hydrogen but one of R^2 and R^4 must be hydrogen; or
 a pharmaceutically acceptable salt thereof.

9. The ^{225}Ac conjugate of Claim 6 wherein the
 functionalized chelant is a compound of formula IV



wherein:

- 5 each Q independently is hydrogen or CHR^5COOR ; with the proviso that at least one Q must be other than hydrogen;
- each R independently is hydrogen, benzyl or $\text{C}_1\text{-C}_4$ alkyl;
- 10 m is integer from 0 to 5 inclusive;
- R^2 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 15 R^3 is selected from the group consisting of $\text{C}_1\text{-C}_4$ alkoxy, $-\text{OCH}_2\text{COOH}$, hydroxy and hydrogen;
- R^4 is selected from the group consisting of hydrogen, nitro, amino, isothiocyanato, semicarbazido, thiosemicarbazido, carboxyl, bromoacetamido and maleimido;
- 20 each R^5 independently is hydrogen or $\text{C}_1\text{-C}_4$ alkyl; with the proviso that R^2 and R^4 cannot both be hydrogen but one of R^2 and R^4 must be hydrogen; or a pharmaceutically acceptable salt thereof.

25

10. The ^{225}Ac conjugate of Claim 6 wherein the functionalized chelant compound is 1-[(2-methoxy-5-isothiocyanatophenyl)-carboxymethyl]-4,7,10-triscarboxymethyl-1,4,7,10-tetraazacyclododecane (MeO-DOTA-NCS).

30

11. The ^{225}Ac conjugate of any one of Claims 6 to 10 wherein the biological molecule is an antibody or antibody fragment.

5 12. The ^{225}Ac conjugate of any one of Claims 6 to 10 wherein the biological molecule is selected from the group of antibodies consisting of NuM195, CC-49, CC-49 F(ab')₂, CC-83, and CC-83 F(ab')₂.

10 13. The ^{225}Ac conjugate of Claim 6 wherein the functionalized chelant compound of the conjugate is 1-[(2-methoxy-5-isothiocyanatophenyl)-carboxymethyl]-4,7,10-triscarboxy-methyl-1,4,7,10-tetraazacyclododecane and the
15 biological molecule is selected from the group of antibodies consisting of HuM195, CC-49, CC-49 F(ab')₂, CC-83, and CC-83 F(ab')₂.

14. The ^{225}Ac conjugate of Claim 13 wherein the functionalized chelant compound of the conjugate is 1-[(2-
20 methoxy-5-isothiocyanatophenyl)-carboxymethyl]-4,7,10-triscarboxy-methyl-1,4,7,10-tetraazacyclododecane and the biological molecule is HuM195 antibody.

15 15. A pharmaceutical formulation comprising the ^{225}Ac conjugate of any one of Claims 6 to 10 with a pharmaceutically acceptable carrier.

16. A pharmaceutical formulation comprising the ^{225}Ac conjugate of Claim 12 with a pharmaceutically acceptable
30 carrier.

17. A pharmaceutical formulation comprising the ^{225}Ac conjugate of Claim 13 or Claim 14 with a pharmaceutically acceptable carrier.

35

18. A method of the therapeutic treatment of a mammal having cancer which comprises administering to said mammal a therapeutically effective amount of a pharmaceutical formulation of Claim 15.

5

19. A method of the therapeutic treatment of a mammal having cancer which comprises administering to said mammal a therapeutically effective amount of a pharmaceutical formulation of Claim 16.

10

20. A method of the therapeutic treatment of a mammal having cancer which comprises administering to said mammal a therapeutically effective amount of a pharmaceutical formulation of Claim 17.